




**Environmental  
Product  
Declaration**

According to EN15804+A2 (+indicators A1)



This declaration is for:  
**DF2**

Provided by:  
**Ankersmit Maalbedrijven B.V.**



program operator  
**Stichting MRPI®**  
publisher  
**Stichting MRPI®**  
[www.mrpi.nl](http://www.mrpi.nl)

MRPI® registration  
**1.1.00533.2024**  
date of first issue  
**1-5-2024**  
date of this issue  
**1-5-2024**  
expiry date  
**1-5-2029**



**COMPANY INFORMATION**



Ankersmit Maalbedrijven B.V.  
Op de Bos 300  
6223 EP MAASTRICHT  
043 – 366 3737  
[Customersupport.nl@sibelco.com](mailto:Customersupport.nl@sibelco.com)  
<https://www.sibelco.com>

**PRODUCT**

DF2

**DECLARED UNIT/FUNCTIONAL UNIT**

1000 kg

**DESCRIPTION OF PRODUCT**

Composite filler. Intended use according to DoP: Fillers for asphalt and surface treatment for roads, airfields and other traffic areas.

**MRPI® REGISTRATION**

1.1.00533.2024


**DATE OF ISSUE**

1-5-2024

**EXPIRY DATE**

1-5-2029

**VISUAL PRODUCT**




**SCOPE OF DECLARATION**

This MRPI®-EPD certificate is verified by Anne Kees Jeeninga, Advieslab VOF. The LCA study has been done by Bob Roijen, SGS INTRON. The certificate is based on an LCA-dossier according to EN15804+A2 (+indicators A1). It is verified according to the 'MRPI®-EPD verification protocol November 2020.v4.0'. EPD's of construction products may not be comparable if they do not comply with EN15804+A2. Declaration of SVHC that are listed on the 'Candidate list of Substances of Very High Concern for authorisation' when content exceeds the limits for registration with ECHA.

**MORE INFORMATION**

**PROGRAM OPERATOR**

Stichting MRPI®  
Kingsfordweg 151  
1043 GR  
Amsterdam

DEMONSTRATION OF VERIFICATION	
CEN standard EN15804 serves as the core PCR(a)	
Independent verification of the declaration an data according to EN15804+A2 (+indicators A1)	
internal:	external: x
Third party verifier: Anne Kees Jeeninga, Advieslab VOF	
	
[a] PCR = Product Category Rules	

Ing. L. L. Oosterveen MSc. MBA  
Managing Director MRPI

### DETAILED PRODUCT DESCRIPTION


Raw materials are ground and mixed in ball mills. Damp materials are dried in order to yield a dry end-product. This drying process consumes natural gas. Some products are produced by mixing intermediates. All the drying, grinding, and mixing occur at Sibelco's facility. The product is delivered as bulk product.

Component > 1% of total mass	(%)
Raw material	amount
Primary raw materials	approx 30 – 40%
Secondary raw materials	approx 60 – 70%

### SCOPE AND TYPE

The filler is produced at the location of Ankerpoort Maalbedrijven in Maastricht. Since filler is a semi-finished product only the production phase is included. The LCA calculations have been made in Simapro using Ecoinvent database v3.6 (allocation cut-off) and the NMD process database v3.3.

PRODUCT STAGE		CONSTRUCTION PROCESS STAGE			USER STAGE								END OF LIFE STAGE				BENEFITS AND LOADS BEYOND THE SYSTEM BOUNDARIES
Rawmaterial supply	Transport	Manufacturing	Transport gate to site	Assembly	Use	Maintenance	Repair	Replacement	Refurbishment	Operational energy use	Operational water use	De-construction demolition	Transport	Waste processing	Disposal	Reuse- Recovery – Recycling-potential	
A1	A2	A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D	
X	X	X	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
X= Modules Assessed ND= Not Declared																	



**REPRESENTATIVENESS**

Not applicable, in this study a specific product is considered produced at a specific production site.

**ENVIRONMENT IMPACT per functional unit or declared unit (core indicators A1)**

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
ADPE	kg Sb eq.	3,23 E-06	6,71 E-04	5,15 E-05	7,26 E-04	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ADPF	MJ	2,76 E+00	4,04 E+02	4,68 E+02	8,75 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
GWP	kg CO2 eq.	2,07 E-01	2,65 E+01	3,06 E+01	5,73 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ODP	Kg CFC11 eq.	3,23 E-08	4,72 E-06	3,16 E-06	7,91 E-06	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
POCP	Kg ethene eq.	2,34 E-04	1,60 E-02	5,46 E-03	2,17 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
AP	kg SO2 eq.	1,54 E-03	1,15 E-01	3,61 E-02	1,53 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
EP	kg (PO4) 3- eq.	3,25 E-04	2,26 E-02	6,31 E-03	2,92 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

Toxicity indicators for Dutch market

HTP	kg DCB-Eq	9,60 E-02	1,12 E+01	3,90 E+00	1,52 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
FAETP	kg DCB-Eq	1,18 E-03	3,27 E-01	5,29 E-02	3,81 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
MAETP	kg DCB-Eq	4,45 E+00	1,18 E+03	1,98 E+02	1,38 E+03	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
TETP	kg DCB-Eq	1,61 E-04	3,95 E-02	3,04 E-02	7,01 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ECI	euro	2,93 E-02	3,19 E+00	2,16 E+00	5,38 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ADPF	kg Sb eq.	1,35 E-03	1,95 E-01	2,56 E-01	4,53 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

- ADPE = Abiotic Depletion Potential for non-fossil resources
- ADPF = Abiotic Depletion Potential for fossil resources
- GWP = Global Warming Potential
- ODP = Depletion potential of the stratospheric ozone layer
- POCP = Formation potential of tropospheric ozone photochemical oxidants
- AP = Acidification Potential of land and water
- EP = Eutrophication Potential
- HTP = Human Toxicity Potential
- FAETP = Fresh water aquatic ecotoxicity potential
- MAETP = Marine aquatic ecotoxicity potential
- TETP = Terrestrial ecotoxicity potential
- ECI = Environmental Cost Indicator
- ADPF = Abiotic Depletion Potential for fossil resources expressed in [kg Sb-eq.]



**ENVIRONMENT IMPACT per functional unit or declared unit (core indicators A2)**

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
GWP-total	kg CO2 eq.	2,12 E-01	2,67 E+01	3,12 E+01	5,81 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
GWP-fossil	kg CO2 eq.	2,12 E-01	2,67 E+01	3,10 E+01	5,80 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
GWP-biogenic	kg CO2 eq.	6,15 E-04	1,26 E-02	1,08 E-01	1,21 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
GWP-luluc)	kg CO2 eq.	2,45 E-05	9,74 E-03	4,75 E-03	1,45 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ODP	kg CFC11 eq.	4,07 E-08	5,92 E-06	3,53 E-06	9,49 E-06	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
AP	mol H+ eq.	2,13 E-03	1,53 E-01	4,61 E-02	2,01 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
EP-freshwater	kg PO4 eq.	1,95 E-06	2,68 E-04	5,87 E-04	8,57 E-04	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
EP-marine	kg N eq.	8,64 E-04	5,36 E-02	1,18 E-02	6,63 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
EP-terrestrial	mol N eq.	9,59 E-03	5,91 E-01	1,36 E-01	7,36 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
POCP	kg NMVOC eq.	2,68 E-03	1,69 E-01	4,03 E-02	2,12 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ADP-minerals & metals	kg Sb eq.	3,23 E-06	6,71 E-04	5,15 E-05	7,26 E-04	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ADP-fossil	MJ, net calorific value	2,76 E+00	4,04 E+02	4,68 E+02	8,75 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
WDP	m3 world eq. Deprived	1,50 E-02	1,44 E+00	1,27 E+00	2,72 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

- GWP-total = Global Warming Potential total
- GWP-fossil = Global Warming Potential fossil fuels
- GWP-biogenic = Global Warming Potential biogenic
- GWP-luluc = Global Warming Potential land use and land use change
- ODP = Depletion potential of the stratospheric ozone layer
- AP = Acidification Potential, Accumulated Exceedence
- EP-freshwater = Eutrophication Potential, fraction of nutrients reaching freshwater end compartment
- EP-marine = Eutrophication Potential, fraction of nutrients reaching marine end compartment
- EP-terrestrial = Eutrophication Potential, Accumulated Exceedence
- POCP = Formation potential of tropospheric ozone photochemical oxidants
- ADP-minerals&metals = Abiotic Depletion Potential for non-fossil resources [2]
- ADP-fossil = Abiotic Depletion for fossil resources potential [2]
- WDP = Water (user) deprivation potential, deprivation-weighted water consumption [2]

Disclaimer [2]

- The results of this environmental impact indicator shall be used with care as the uncertainties on these results are high or as there is limited experienced with the indicator



**ENVIRONMENT IMPACT per functional unit or declared unit (additional indicators A2)**

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
PM	Disease incidence	6,05 E-08	2,40 E-06	2,89 E-07	2,75 E-06	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
IRP	kBq U235 eq.	1,12 E-02	1,69 E+00	4,31 E-01	2,14 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ETP- fw	CTUe	2,66 E+00	3,59 E+02	1,72 E+02	5,33 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
HTP- c	CTUh	7,76 E-11	1,16 E-08	7,95 E-09	1,96 E-08	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
HTP- nc	CTUh	3,48 E-09	3,93 E-07	1,49 E-07	5,45 E-07	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
SQP	----	3,93 E+00	3,54 E+02	3,19 E+01	3,90 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

PM = Potential incidence of disease due to PM emissions

IRP = Potential Human exposure efficiency relative to U235 [1]

ETP-fw = Potential Comparative Toxic Unit for ecosystems [2]

HTP-c = Potential Comparative Toxic Unit for humans [2]

HTP-nc = Potential Comparative Toxic Unit for humans, non-cancer [2]

SQP = Potential soil quality index [2]

**Disclaimer [1]**

- This impact category deals mainly with the eventual impact of low dose ionizing radiation on human health of the nuclear fuel cycle. It does not consider effects due to possible nuclear accidents, occupational exposure nor due to radioactive waste.

**Disclaimer [2]**

- The results of this environmental impact indicator shall be used with care as the uncertainties on these results are high or as there is limited experienced with the indicator.

**OUTPUT FLOWS AND WASTE CATEGORIES per functional unit or declared unit (A1 / A2)**

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
HWD	kg	7,57 E-06	1,02 E-03	2,22 E-02	2,32 E-02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
NHWD	kg	9,78 E-03	2,59 E+01	1,37 E+00	2,73 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
RWD	kg	1,80 E-05	2,66 E-03	5,11 E-04	3,18 E-03	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
CRU	kg	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
MFR	kg	0,00 E+00	0,00 E+00	1,43 E-01	1,43 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
MER	kg	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
EEE	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
ETE	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

HWD = Hazardous Waste Disposed  
 RWD = Radioactive Waste Disposed  
 MFR = Materials for recycling  
 EEE = Exported Electrical Energy  
 NHWD = Non Hazardous Waste Disposed  
 CRU = Components for reuse  
 MER = Materials for energy recovery  
 ETE = Exported Thermal Energy



### RESOURCE USE per functional unit or declared unit (A1 / A2)

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
PERE	MJ	4,05 E-02	5,06 E+00	1,30 E+01	1,81 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PERM	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PERT	MJ	4,05 E-02	5,06 E+00	1,30 E+01	1,81 E+01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PENRE	MJ	2,93 E+00	4,29 E+02	5,14 E+02	9,46 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PENRM	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
PENRT	MJ	2,93 E+00	4,29 E+02	5,14 E+02	9,46 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
SM	kg	6,63 E+02	0,00 E+00	0,00 E+00	6,63 E+02	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
RSF	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
NRSF	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
FW	m3	7,42 E-03	4,91 E-02	8,62 E-02	1,43 E-01	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

PERE = Use of renewable energy excluding renewable primary energy resources

PERM = Use of renewable energy resources used as raw materials

PERT = Total use of renewable primary energy resources

PENRE = Use of non-renewable primary energy resources excluding non-renewable energy resources used as raw materials

PENRM = Use of non-renewable primary energy resources used as raw materials

PENRT = Total use of non-renewable primary energy resources

SM = Use of secondary materials

RSF = Use of renewable secondary fuels

NRSF = Use of non-renewable secondary fuels

FW = Use of net fresh water

### BIOGEEEN CARBON CONTENT per functional unit or declared unit (A1 / A2)

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	B3	B4	B5	B6	B7	C1	C2	C3	C4	D
BBCpr	Kg C	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
BCCpa	kg C	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

BBCpr = Biogenic carbon content in product

BCCpa = Biogenic carbon content in packaging



### **CALCULATION RULES**

This EPD is based on production data collected in 2021 over a complete production year: 2020. Virtually no materials or processes have been excluded from the study (cut-off rule is well below 1%). In case secondary materials are applied transport to the production location of Sibelco and processing (eg: drying) is allocated to the production of the filler. Infrastructure processes in Ecoinvent processes have been included, long term emissions in Ecoinvent processes have been excluded from the LCA calculations.

### **SENARIOS AND ADDITIONAL TECHNICAL INFORMATION**

The filler is produced by milling and grinding raw materials in crushers and ball mills. Moisture in raw materials is evaporated in order to yield a dry end product. The drying process consumes natural gas. Some of the products are produced by mixing intermediate products. However all of the drying, grinding and mixing takes place at the facilities of Sibelco and is included in the data collection and LCA calculations.

### **DECLARATION OF SVHC**

No substances that are listed in the latest "Candidate List of Substances of Very High Concern for authorisation" are included in the product that exceeds the limit for registration.

### **REFERENCES**

- Stichting Nationale Milieudatabase, Bepalingsmethode Milieuprestatie Bouwwerken Versie 1.0 (juli 2020).
- EN 15804:2012+A2:2019, Sustainability of construction works - Environmental product declarations - Core rules for the product category of construction products, 2019.
- ISO 14040:2006. Environmental management — Life cycle assessment — Principles and framework. 2006.
- ISO 14044:2006. Environmental management — Life cycle assessment — Requirements and guidelines. 2006.
- ISO 21930:2017. Sustainability in buildings and civil engineering works — Core rules for environmental product declarations of construction products and services. 2017.
- ISO 14025:2006. Environmental labels and declarations — Type III environmental declarations — Principles and procedures. 2006.
- SGS INTRON report: A117300/R20210185b, January 6th 2022
- DoP 0431/004 DF2 modifications are possible to a limited extent.

### **REMARKS**

None.