



**Environmental** 

**Product** 

**Declaration** 

According to EN15804+A2 (+indicators A1)



This declaration is for:

**M6** 

Provided by:

Sibelco Benelux BV





program operator
Stichting MRPI®
publisher
Stichting MRPI®
www.mrpi.nl

MRPI® registration
1.1.00536.2024
date of first issue
31-5-2024
date of this issue
31-5-2024
expiry date
31-5-2029











#### **COMPANY INFORMATION**



Sibelco Benelux BV
Tiegelstraat 1
6414 XB HEERLEN
045 – 563 6414
customersupport.nl@sibelco.com
https://www.sibelco.com/en



#### **MRPI® REGISTRATION**

1.1.00536.2024

**DATE OF ISSUE** 

31-5-2024

**EXPIRY DATE** 

31-5-2029



#### **SCOPE OF DECLARATION**

This MRPI®-EPD certificate is verified by Anne Kees Jeeninga, Advieslab VOF. The LCA study has been done by Bob Roijen, SGS INTRON. The certificate is based on an LCA-dossier according to EN15804+A2 (+indicators A1). It is verified according to the 'MRPI®-EPD verification protocol November 2020.v4.0'. EPD's of construction products may not be comparable if they do not comply with EN15804+A2. Declaration of SVHC that are listed on the 'Candidate list of Substances of Very High Concern for authorisation' when content exceeds the limits for registration with ECHA.



#### **PRODUCT**

M6



## **DECLARED UNIT/FUNCTIONAL UNIT**

1000 kg



#### **DESCRIPTION OF PRODUCT**

Filler, natural silica. Intended use according to DoP: fillers for concrete and fillers for asphalt and surface treatment for roads, airfields and other traffic areas.



#### **VISUAL PRODUCT**



**MORE INFORMATION** 



# **PROGRAM OPERATOR**

Stichting MRPI®
Kingsfordweg 151
1043 GR
Amsterdam



Lw Cookwen

Ing. L. L. Oosterveen MSc. MBA Managing Director MRPI

## **DEMONSTRATION OF VERIFICATION**

CEN standard EN15804 serves as the core PCR(a)

Independent verification of the declaration an data according to EN15804+A2 (+indicators A1)

internal:

external: x

Third party verifier: Anne Kees Jeeninga, Advieslab VOF

[a] PCR = Product Category Rules









## **DETAILED PRODUCT DESCRIPTION**

Raw materials are ground and mixed in ball mills. Damp materials are dried in order to yield a dry end-product. This drying process consumes natural gas. Some products are produced by mixing intermediates. All the drying, grinding, and mixing occur at Sibelco's facility.

The product is delivered as bulk product.



Component > 1% of total mass	(%)
Raw material	Amount
Silica	100%



#### **SCOPE AND TYPE**

The filler is produced at the location of Sibelco Benelux BV in Heerlen. Since filler is a semi-finished product only the production phase is included. The LCA calculations have been made in Simapro using Ecoinvent database v3.6 (allocation cut-off) and the NMD process database v3.3.



			ltion ltion											THE SYSTEM BOUNDARIES
Rawmaterial supply Transport Manufacturing	Transport gate to site	Assembly	Use	Maintenance	Repair	Replacement	Refurbishment	Operational energy use	Operational water use	De-construction demolition	Transport	Waste processing	Disposal	Reuse- Recovery – Recycling- potential
A1 A2 A3	A4	A5	A5 B1 B2 B3 B4 B5 B6 B7 C1 C2 C3						C3	C4	D			
X X X	ND	ND	ID ND					ND	ND					


X= Modules Assessed

ND= Not Declared











## **REPRESENTATIVENESS**

Not applicable, in this study a specific product is considered produced at a specific production site.







# **ENVIRONMENT IMPACT per functional unit or declared unit (core indicators A1)**

	Unit	A1	A2	А3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	СЗ	C4	D
ADPE	kg Sb eq.	1,11 E-06	2,40 E-06	8,87 E-05	9,22 E-05	ND													
ADPF	MJ	9,94 E+00	1,43 E+00	4,65 E+02	4,76 E+02	ND													
GWP	kg CO2 eq.	7,15 E-01	9,41 E-02	3,40 E+01	3,48 E+01	ND													
ODP	Kg CFC11 eq.	1,24 E-07	1,67. E-08	2,10 E-06	2,24 E-06	ND													
РОСР	Kg ethene eq.	7,28 E-04	5,68 E-05	5,55 E-03	6,33 E-03	ND													
AP	kg SO2 eq.	5,39 E-03	4,14 E-04	5,85 E-02	6,43 E-02	ND													
EP	kg (PO4 ) 3- eq.	1,22 E-03	8,13 E-05	1,19 E-02	1,32 E-02	ND													
Toxicity ind	licators for Di	utch market																	
НТР	kg DCB-Eq	2,65 E-01	3,96 E-02	4,64 E+00	4,95 E+00	ND													
FAETP	kg DCB-Eq	3,68 E-03	1,16 E-03	1,10 E-01	1,15 E-01	ND													
MAETP	kg DCB-Eq	1,28 E+01	4,16 E+00	4,19 E+02	4,36 E+02	ND													
TETP	kg DCB-Eq	4,36 E-04	1,40 E-04	6,40 E-02	6,45 E-02	ND													
ECI	euro	9,57 E-02	1,13 E-02	2,56 E+00	2,67 E+00	ND													
ADPF	kg Sb eq.	4,71 E-03	6,92 E-04	2,61 E-01	2,67 E-01	ND													

ADPE = Abiotic Depletion Potential for non-fossil resources

ADPF = Abiotic Depletion Potential for fossil resources

GWP = Global Warming Potential

ODP = Depletion potential of the stratospheric ozone layer

POCP = Formation potential of tropospheric ozone photochemical oxidants

AP = Acidification Potential of land and water

EP = Eutrophication Potential HTP = Human Toxicity Potential

FAETP = Fresh water aquatic ecotoxicity potential
MAETP = Marine aquatic ecotoxicity potential
TETP = Terrestrial ecotoxicity potential

ECI = Environmental Cost Indicator

ADPF = Abiotic Depletion Potential for fossil resources expressed in [kg Sb-eq.]









# **ENVIRONMENT IMPACT** per functional unit or declared unit (core indicators A2)

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	C3	C4	D
GWP-	kg	7,22	9,50	3,47	3,56														
total	CO2 eq.	E-01	E-02	E+01	E+01	ND													
GWP-	kg	7,22	9,49	3,44	3,52														
fossil	CO2 eq.	E-01	E-02	E+01	E+01	ND													
GWP-	kg	2,01	4,38	3,20	3,21														
biogenic	CO2 eq.	E-04	E-05	E-01	E-01	ND													
GWP-	kg	5,69	3,48	1,01	1,01														
luluc)	CO2 eq.	E-05	E-05	E-02	E-02	ND													
ODP	kg	1,56	2,09	2,17	2,35														
ODI	CFC11 eq.	E-07	E-08	E-06	E-06	ND													
AP	mol	7,55	5,50	7,43	8,24														
All	H+ eq.	E-03	E-04	E-02	E-02	ND													
EP-	kg	2,63	9,57	1,68	1,68														
freshwater	PO4 eq.	E-06	E-07	E-03	E-03	ND													
EP-	kg	3,33	1,94	1,67	2,02														
marine	N eq.	E-03	E-04	E-02	E-02	ND													
EP-	mol	3,66	2,14	2,01	2,40														
terrestrial	N eq.	E-02	E-03	E-01	E-01	ND													
POCP	kg	1,01	6,10	5,19	6,26														
1 001	NMVOC eq.	E-02	E-04	E-02	E-02	ND													
ADP-minerals	kg	1,11	2,40	8,87	9,22														
& metals	Sb eq.	E-06	E-06	E-05	E-05	ND													
ADP-fossil	MJ, net	9,94	1,43	4,65	4,76														
103311	calorific value	E+00	E+00	E+02	E+02	ND													
WDP	m3 world	1,33	5,12	3,26	3,27														
	eq. Deprived	E-02	E-03	E+00	E+00	ND													

GWP-total = Global Warming Potential total

GWP-fossil = Global Warming Potential fossil fuels

GWP-biogenic = Global Warming Potential biogenic

GWP-luluc = Global Warming Potential land use and land use change

 $\label{eq:ode_potential} \textsc{ODP} = \textsc{Depletion potential of the stratospheric ozone layer}$ 

AP = Acidification Potential, Accumulated Exceedence

EP-freshwater = Eutrophication Potential, fraction of nutrients reaching freshwater end compartment

EP-marine = Eutrophication Potential, fraction of nutrients reaching marine end compartment

EP-terrestrial = Eutrophication Potential, Accumulated Exceedence

 ${\tt POCP = Formation\ potential\ of\ tropospheric\ ozone\ photochemical\ oxidants}$ 

ADP-minerals&metals = Abiotic Depletion Potential for non-fossil resources [2]

ADP-fossil = Abiotic Depletion for fossil resources potential [2]

WDP = Water (user) deprivation potential, deprivation-weighted water consumption [2]

#### Disclamer [2]

- The results of this environmental impact indicator shall be used with care as the uncertainties on these results are high or as there is limited experienced with the indicator









# **ENVIRONMENT IMPACT per functional unit or declared unit (additional indicators A2)**

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	C3	C4	D
PM	Disease	2,00	8,52	2,78	4,86														
	incidence	E-07	E-09	E-07	E-07	ND													
IRP	kBq	4,26	6,00	8,48	8,97														
	U235 eq.	E-02	E-03	E-01	E-01	ND													
ETP-	CTUe	5,99	1,28	4,15	4,22														
fw	CIOC	E+00	E+00	E+02	E+02	ND													
HTP-	CTUh	2,09	4,14	1,43	1,45														
С	CTOIL	E-10	E-11	E-08	E-08	ND													
HTP-	CTUh	5,14	1,40	3,29	3,36														
nc		E-09	E-09	E-07	E-07	ND													
SQP		1,27	1,24	8,23	8,48														
•		E+00	E+00	E+01	E+01	ND													

PM = Potential incidence of disease due to PM emissions

IRP = Potential Human exposure efficiency relative to U235 [1]

ETP-fw = Potential Comparative Toxic Unit for ecosystems [2]

HTP-c = Potential Comparative Toxic Unit for humans [2]

HTP-nc = Potential Comparative Toxic Unit for humans, non-cancer [2]

SQP = Potential soil quality index [2]

#### Disclamer [1]

- This impact category deals mainly with the eventual impact of low dose ionizing radiation on human health of the nuclear fuel cycle. It does not consider effects due to possible nuclear accidents, occupational exposure nor due to radioactive waste.

## Disclamer [2]

- The results of this environmental impact indicator shall be used with care as the uncertainties on these results are high or as there is limited experienced with the indicator.









# OUTPUT FLOWS AND WASTE CATEGORIES per functional unit or declared unit (A1 / A2)

	Unit	A1	A2	A3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	C3	C4	D
HWD	kα	2,71	3,63	2,20	2,20														
TIVVD	kg	E-05	E-06	E-02	E-02	ND													
NHWD	kg	1,18	9,08	2,88	2,98														
INTIVVD	^g	E-02	E-02	E+00	E+00	ND													
RWD	kg	6,90	9,40	8,66	9,44														
KWD	Ng	E-05	E-06	E-04	E-04	ND													
CRU	kg	0,00	0,00	0,00	0,00														
CKO	∧g	E+00	E+00	E+00	E+00	ND													
MFR	kg	0,00	0,00	1,43	1,43														
IVIIIX	Ng.	E+00	E+00	E-01	E-01	ND													
MER	kg	0,00	0,00	0,00	0,00														
IVILIX	Ng.	E+00	E+00	E+00	E+00	ND													
EEE	MJ	0,00	0,00	0,00	0,00														
	IVIJ	E+00	E+00	E+00	E+00	ND													
ETE	MJ	0,00	0,00	0,00	0,00	,		,	,								,		
LIL	IVIJ	E+00	E+00	E+00	E+00	ND													

HWD = Hazardous Waste Disposed

RWD = Radioactive Waste Disposed

MFR = Materials for recycling

EEE = Exported Electrical Energy

NHWD = Non Hazardous Waste Disposed

CRU = Components for reuse

MER = Materials for energy recovery

ETE = Exported Thermal Energy









# RESOURCE USE per functional unit or declared unit (A1 / A2)

	Unit	A1	A2	А3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	C3	C4	D
PERE	MJ	5,38 E-02	1,79 E-02	3,72 E+01	3,73 E+01	ND													
PERM	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND													
PERT	MJ	5,38 E-02	1,79 E-02	3,72 E+01	3,73 E+01	ND													
PENRE	MJ	1,06 E+01	1,52 E+00	5,00 E+02	5,13 E+02	ND													
PENRM	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND													
PENRT	MJ	1,06 E+01	1,52 E+00	5,00 E+02	5,13 E+02	ND													
SM	kg	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND													
RSF	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND													
NRSF	MJ	0,00 E+00	0,00 E+00	0,00 E+00	0,00 E+00	ND													
FW	m3	5,12 E-04	1,74 E-04	2,43 E-01	2,43 E-01	ND													

PERE = Use of renewable energy excluding renewable primary energy resources

PERM = Use of renewable energy resources used as raw materials

PERT = Total use of renewable primary energy resources

PENRE = Use of non-renewable primary energy resources excluding non-renewable energy resources used as raw materials

PENRM = Use of non-renewable primary energy resources used as raw materials

PENRT = Total use of non-renewable primary energy resources

SM = Use of secondary materials

RSF = Use of renewable secondary fuels

NRSF = Use of non-renewable secondary fuels

FW = Use of net fresh water



# BIOGEEN CARBON CONTENT per functional unit or declared unit (A1 / A2)

	Unit	A1	A2	А3	A1-A3	A4	A5	B1	B2	В3	B4	B5	В6	В7	C1	C2	C3	C4	D
BBCpr	Kg C	0,00	0,00	0,00	0,00														
вьсрі	Ng C	E+00	E+00	E+00	E+00	ND													
ВССра	kg C	0,00	0,00	0,00	0,00														
ВССРА	Ng C	E+00	E+00	E+00	E+00	ND													

BCCpr = Biogenic carbon content in product
BCCpa = Biogenic carbon content in packaging









#### **CALCULATION RULES**

This EPD is based on production data collected in 2021 over a complete production year: 2020. Virtually no materials or processes have been excluded from the study (cut-off rule is well below 1%). In case secondary materials are applied transport to the production location of Sibelco and processing (eg: drying) is allocated to the production of the filler. Infrastructure processes in Ecoinvent processes have been included, long term emissions in Ecoinvent processes have been excluded from the LCA calculations.



#### **SENARIOS AND ADDITIONAL TECHNICAL INFORMATION**

The filler is produced by milling and grinding raw materials in crushers and ball mills. Moisture in raw materials is evaporated in order to yield a dry end product. The drying process consumes natural gas. Some of the products are produced by mixing intermediate products. However all of the drying, grinding and mixing takes place at the facilities of Sibelco and is included in the data collection and LCA calculations.



#### **DECLARATION OF SVHC**

No substances that are listed in the latest "Candidate List of Substances of Very High Concern for authorisation" are included in the product that exceeds the limit for registration.



#### **REFERENCES**

- •ISO 14044:2006. Environmental management Life cycle assessment Requirements and guidelines. 2006.
- •ISO 21930:2017. Sustainability in buildings and civil engineering works Core rules for environmental product declarations of construction products and services. 2017.
- ISO 14025:2006. Environmental labels and declarations Type III environmental declarations Principles and procedures. 2006.
- •SGS INTRON report: A117300/R20210185b, January 6th 2022
- •DoP ADH-M6-2020-01 modifications are possible to a limited extent.
- •DoP DH-M6-2021-12-28 modifications are possible to a limited extent.



#### **REMARKS**

None.

